

COVID-19 in Children: Initial Characterization of the Pediatric Disease

DOI: 10.1542/peds.2020-0834

Journal: *Pediatrics*

Citation: Cruz A, Zeichner S. COVID-19 in children: initial characterization of the pediatric disease. *Pediatrics*. 2020; doi: 10.1542/peds.2020-0834

This is a pre-publication version of an article that has undergone peer review and been accepted for publication but is not the final version of record. This article will be copyedited and typeset before it is published in its final form. The journal is providing an early version of this article to expedite access to these findings.

COVID-19 in Children: Initial Characterization of the Pediatric Disease

Andrea T. Cruz, MD, MPH¹ and Steven L. Zeichner, MD, PhD²

Affiliations: ¹Department of Pediatrics, Baylor College of Medicine, Houston, TX;

²Departments of Pediatrics & Microbiology, Immunology, and Cancer Biology, University of Virginia, Charlottesville, VA

Address correspondence to: Andrea Cruz, Department of Pediatrics, 6621 Fannin Street, Suite A2210, Houston, TX 77030, acruz@bcm.edu, 832-824-5582

Short title: Children and COVID-19

Funding source: no funding was secured for this study.

Financial disclosure: Andrea Cruz and Steven Zeichner are associate editors for *Pediatrics*.

Potential conflicts of interest: Dr. Cruz has no conflict of interest to disclose. The University of Virginia has filed patent applications related to new technologies for the rapid production of vaccines and Dr. Zeichner is an inventor on those patent applications.

Abbreviations: ARDS: acute respiratory distress syndrome; COVID-19: coronavirus disease; SARS-CoV-2: severe acute respiratory syndrome coronavirus 2

The impact of the disease caused by the novel coronavirus, SARS-CoV-2,¹ COVID-19, has been widespread, with over 120,000 cases diagnosed in more than 100 countries since the virus was identified in January of 2020.² Preliminary data focused on severe respiratory manifestations, seen predominantly in adults, with scant initial data on the burden of COVID-19 in children.³ We therefore read with interest the findings of Dong and colleagues in this volume of *Pediatrics*, who reported a series of over 2000 children with suspected or confirmed COVID-19.⁴ The authors found that 13% of virologically-confirmed cases had asymptomatic infection, a rate that almost certainly understates the true rate of asymptomatic infection, since many asymptomatic children are unlikely to be tested. Among symptomatic children, 5% had dyspnea or hypoxemia (a substantially lower percentage than what has been reported for adults³) and 0.6% progressed to acute respiratory distress syndrome (ARDS) or multiorgan system dysfunction, a rate that is also lower than that seen in adults. Preschool-aged children and infants were more likely to have severe clinical manifestations than older children.

There are several salient points from this paper. First, while children are less likely to become severely ill than older adults, there are subpopulations of children with an increased risk for more significant illness. These are consistent with data on non-COVID-19 coronaviruses. One viral surveillance study in a pediatric intensive care unit in China reported that coronavirus was detected in more children with ARDS than human metapneumovirus.⁵ Another study in hospitalized Norwegian children detected coronaviruses in 10% of hospitalized children with respiratory tract infections.⁶ Younger age, underlying pulmonary pathology, and immunocompromising conditions have been associated with more severe outcomes with non-COVID-19 coronavirus infections in children.⁷

Second, the attributable risk for severe disease from COVID-19 in children is challenging to discern. Prior studies have shown that children from whom coronaviruses are detected from the respiratory tract can have viral co-infections in up to two-thirds of cases.⁶ In the study by Dong et al., testing for other viruses was not standardized, and two-thirds of cases were clinically diagnosed, not virologically confirmed. Furthermore, children without virologic confirmation were more likely to have severe disease than children from whom COVID-19 was detected, potentially because their symptoms were caused by other pathogens.

Third, children may play a major role in community-based viral transmission. Available data suggest that children may have more upper respiratory tract (including nasopharyngeal carriage), rather than lower respiratory tract involvement.⁴ There is also evidence of fecal shedding in the stool for several weeks after diagnosis,⁸ leading to concern about fecal-oral transmission of the virus, particularly for infants and children who are not toilet-trained, and for viral replication in the gastrointestinal tract.⁹ Prolonged shedding in nasal secretions and stool has substantial implications for community spread in daycare centers, schools, and in the home. Additionally, non-COVID-19 coronaviruses are detectable in respiratory secretions in a large percentage of healthy children,⁶ and the extent to which this is also seen in COVID-19 is unclear. Prolonged viral shedding in symptomatic individuals, combined with shedding in asymptomatic persons, would render contact tracing and other public health measures to mitigate spread less effective.

We have learned an amazing amount about COVID-19 in a short amount of time, with copious epidemiologic, virologic, and clinical data being published. The SARS-CoV-2 sequence, now published,¹⁰ was first posted to the bioRxiv preprint server a remarkable six weeks after the start

of epidemic, enabling the essential work of molecular epidemiology. The transmission of data has been surpassed only by the transmission of the virus itself. However, there is still much that we need to learn about the impact of this virus on children, as well as the impact of children on viral spread. While vertical transmission has not yet been reported,¹¹ many of the infants born to COVID-19-infected mothers were delivered surgically and quickly separated from their mothers. Many infectious diseases affect pregnant women more severely, and respiratory disease in pregnant women may result in poor fetal outcomes. Data on the basic reproductive number of the virus (the number of persons to whom an infected individual transmits the virus) have varied widely,^{12,13} and household studies can refine data we have on viral transmission and on viral shedding. Widespread availability of testing will allow for us to more accurately describe the spectrum of illness and may result in adjustment of the apparent morbidity and mortality rate as fewer ill individuals are diagnosed. While the focus on pandemics often is on the impact on the persons who utilize the highest resources or on the economically productive age groups, rigorously gauging the impact of COVID-19 on children will be important to accurately model the pandemic and to ensure that appropriate resources are allocated to children requiring care. Many infectious diseases affect children differently than adults and understanding those differences can yield important insights into disease pathogenesis, informing management and the development of therapeutics. This will likely be true for COVID-19, just as it was for older infectious diseases.

References

1. Gorbalenya AE, Baker SC, Baric RS, et al. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat Microbiol.* 2020. doi: 10.1038/s41564-020-0695-z.
2. Johns Hopkins University Center for Systems Science and Engineering. Coronavirus COVID-19 global cases. Available online at: <https://gisanddata.maps.arcgis.com/apps/opsdashboard/index.html#/bda7594740fd40299423467b48e9ecf6>. Last accessed March 11, 2020.
3. Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA.* 2020; doi: 10.1001/jama.2020.1585.
4. Dong Y, Mo X, Hu, Y, et al. [Epidemiology of COVID-19 among children in China]. IN PRESS this issue *Pediatrics*.
5. Li YT, Liang Y, Ling YS, et al. The spectrum of viral pathogens in children with severe acute lower respiratory tract infection: a 3-year prospective study in the pediatric intensive care unit. *J Med Virol.* 2019;91(9): 1633-1642.
6. Heimdal I, Moe N, Kokstad S, et al. Human coronavirus in hospitalized children with respiratory tract infections: a 9-year population-based study from Norway. *J Infect Dis.* 2019;219(8):1198-1206.
7. Omigi C, Englund JA, Bradford MC, Qin X, Boeckh M, Waghmare A. Characteristics and outcomes of coronavirus infection in children: role of viral factors and an immunocompromised state. *J Pediatr Infect Dis Soc.* 2019;8(1):21-28.
8. Jiehao C, Jing X, Daojiong L, et al. A case series of children with 2019 novel coronavirus infection: clinical and epidemiological features. *Clin Infect Dis.* 2020; doi: 10.1093/cid/ciaa198.
9. Xiao F, Tang M, Zheng X, Liu Y, Li X, Shan H. Evidence for gastrointestinal infection of SARS-CoV-2. *Gastroenterol.* 2020. doi: 10.1053/j.gastro.2020.02.055.
10. Zhou P, Yang XL, Wang XG, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature.* 2020. doi: 10.1038/s41586-020-201207.
11. Chen H, Guo J, Wang C, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet.* 2020;395(10226):809-815.
12. Jung SM, Akhmetzhanov AR, Hayashi K, et al. Real-time estimation of the risk of death from novel Coronavirus (COVID-19) infection: inference using exported cases. *J Clin Med.* 2020;9(2): doi: 10.3390/jcm9020523.
13. Hellewell J, Abbott S, Gimma A, et al. Feasibility of controlling COVID-19 outbreaks by isolation of cases and controls. *Lancet Glob Health.* 2020; doi: 10.1016/S2214-109X(20)30074-7.

COVID-19 in Children: Initial Characterization of the Pediatric Disease
Pediatrics originally published online March 16, 2020;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/early/2020/03/16/peds.2020-0834.citation
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Adolescent Health/Medicine http://www.aappublications.org/cgi/collection/adolescent_health:medicine_sub Infectious Disease http://www.aappublications.org/cgi/collection/infectious_diseases_sub HIV/AIDS http://www.aappublications.org/cgi/collection/hiv:aids_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

COVID-19 in Children: Initial Characterization of the Pediatric Disease
Pediatrics originally published online March 16, 2020;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/early/2020/03/16/peds.2020-0834.citation>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2020 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

